



Application of 4,4'-Diaminobenzanilide Schiff Base Metal Complexes As Anti-Tumor and Anti - Dengue Agents

M. Seethalakshmi, T. Peter Amaladhas*

Department of Chemistry, V.O.Chidambaram College, Tuticorin, affiliated to Manonmaniam Sundaranar University, Tirunelveli, TN, India.

Abstract

A unique Schiff base ligand 4-(4-nitrobenzylideneamino)-N-(4-(4-nitrobenzylideneamino)phenyl)benzamide L has been prepared by condensing 4,4'-diaminobenzanilide and p-nitrobenzaldehyde. The cobalt(II), nickel(II) and copper(II) complexes of L were also prepared. The formation and the structure of the ligand and its complexes were confirmed by ¹H-NMR, FT-IR, UV-Vis., EPR and EI- mass techniques. The synthesized complexes were docked with Human DNA topoisomerase I (PDB: 1SC7) and Dengue NS3 protease-helicase (PDB ID: 2VBC) using Auto Dock vina and Discovery studio software. The biological applications of the synthesized complexes were carried out by Cytotoxic screening analysis, DNA binding ability by using electronic spectra and Anti-Tumor activity by MTT assay. The results established that the synthesized transition metal complexes can be act as good anti-tumor and anti-dengue agents.

Keywords: Anti-dengue drug; Docking; Schiff base ligand; 4,4'-diaminobenzanilide; p- nitrobenzaldehyde; NS3 protease- helicase.

1. INTRODUCTION

Schiff bases are condensation products of carbonyl compound especially aldehydes or ketones with primary amines and they were first reported by Hugo Schiff [1] in 1864. Several investigations [2-7] on Schiff base showed that the presence of lone pair of electron in nitrogen atom of imine group is of considerable chemical and biological applications. Nowadays, the research field dealing with Schiff base coordination chemistry has expanded enormously. The importance of Schiff base complexes for bioinorganic chemistry, catalysis, biomedical applications, supramolecular chemistry and formation of compounds with unusual properties and structures has been well studied and reviewed [8-13]. Research on interactions between the transition metal complexes (usually Co(II), Ni(II) and Cu(II)) and DNA helps to give a knowledge about development of novel chemotherapeutics and medicine [14-17]. The interactions between small molecules with DNA can often cause DNA damage in cancer cells [18-20]. Dengue virus basically belongs to Flavivirus genus. This virus is a member of Flaviviridae family [21, 22]. The major source of the dengue virus development is human and monkey. Usually dengue affected humans have severe flu, fever, body pain like

symptoms and body temperature almost reaches 40 °C which makes human to be critical. Severe headache, facial flushing and skin rash are extreme symptoms of Dengue fever and this severe condition is said to be Dengue Hemorrhagic Fever (DHF) and Dengue Shock Syndrome (DSS). It is not possible to do the all experiment in low cost. Docking is the only way to give the corresponding exact results of newly synthesized drugs towards the targeting molecule [23-29]. Auto dock vina and discovery studio is a useful tool to study the interaction between the ligand and the target molecule. This software helps to determine the binding mode, binding affinity and also the exact stable configuration of the ligand which binds with the receptor [30-33]. Using this docking study, we can understand the coordination mode and the binding nature. It will help to design a good drug for the disease [34-37]. In this present work we plan to synthesize novel Schiff base ligand (L) and its cobalt(II), nickel(II) and copper(II) metal complexes, which were treated with human DNA topoisomerase I (PDB: 1SC7) and Dengue NS3 protease-helicase bi-functional enzyme (PDB ID: 2VBC) by docking. The biological applications like cytotoxic screening, DNA binding ability using electronic spectral studies and anti-tumor activity using MTT assay.

*Peter Amaladhas

email: peteramaladhas@gmail.com

2. MATERIALS & METHODS

All the chemicals used in this work were purchased from commercial sources and used the chemicals without further purification. 4,4'-diaminobenzanilide, *p*-nitrobenzaldehyde, cobalt chloride hexahydrate, nickel chloride hexahydrate and copper chloride dihydrate were bought from Sigma Aldrich, USA and used as received. Solvents used in the present research were purchased from Merck and used without further purification.

¹H-NMR spectrum of the synthesized Schiff base ligands have been recorded in DMSO (d₆) by using TMS as an internal standard on a Bruker Advance DRX 300 FT-NMR instrument. The EI-Mass spectra were recorded using JEOL DX-303 EI mass spectrometer at Indian Institute of Technology, Chennai. Infrared spectra of the solid samples were recorded in JASCO/FT-IR410 spectrometer in the range of 4000 – 400 cm⁻¹ at 16 scans/min. Potassium bromide disc method was employed for sample preparation. Electronic spectra of the complexes were recorded using Perkin Elmer Lambda-25 UV-Vis. spectrophotometer using DMSO as solvent in the range of 200-800 nm. The room temperature X- band EPR spectra of the copper complex in DMSO were recorded on Varian E-4

X-band spectrometer using DPPH as the g-marker at Indian Institute of Technology, Chennai.


EXPERIMENTAL PROCEDURES

Preparation of Schiff base ligand L


A hot solution of 1.136 g (5 mmol) 4,4'-diaminobenzanilide in 20 mL methanol was added slowly to a hot stirring solution of 1.5112 g (10 mmol) *p*-nitrobenzaldehyde in 20 mL methanol. The above mixture was stirred under reflux for 5 hours. On cooling to room temperature, the Schiff bases obtained are filtered, washed with diethyl ether and dried *in vacuo*.

Preparation of Cobalt(II), Nickel(II) and Copper (II) Schiff base (L1) metal complexes

To the hot stirring solution of the 0.9869 g (2 mmol) Schiff base ligand L2 in 20 mL of methanol the corresponding metal(II) chlorides [CoCl₂.6H₂O (0.237 g, 1 mmol), NiCl₂.6H₂O (0.2379 g, 1 mmol) and CuCl₂.2H₂O (0.170 g, 1 mmol)] in 20 mL of methanol were added, stirred under reflux for 6 hours. Then the product obtained was filtered, washed and dried *in vacuo*.



Scheme 1. Preparation of Schiff base Ligand L



Scheme 2. Preparation of Schiff base metal complexes

DNA Binding Studies using electronic absorption spectra

Electronic absorption spectrum of the complex was recorded before and after addition of CT-DNA in the presence of 50 mM Tris-HCl buffer (pH 7.5), Tris-hydrochloride (197 mg, 5mM) and sodium chloride (730 mg, 50 mM) were accurately weighed and made up to 250 mL in a standard measuring flask using double distilled water. The pH of the solution was adjusted to 7.5 using 1 mM sodium hydroxide solution with the help of pH meter before making up to the mark. A fixed concentration of metal complexes (10 μ M) was titrated with incremental amounts of CT-DNA over the range (0 – 200 μ M).

Cytotoxicity screening analysis

The stock culture of bacteria was revived by inoculating in broth medium and grown at 37° C for 18 hrs. The Lysogeny broth (LB) Agar plates were prepared and wells were made in the solidified LB agar plate. Each plate was inoculated with 18 h old cultures (100 μ L, 10⁻⁴ CFU) and spread evenly on the plate. After 20 min, the wells were filled with compound at different concentrations. Standard compound plate was also prepared in the same manner. All the plates were incubated at 37° C for 24 h and the diameter of inhibition zone were noted.

MTT assay

To determine the cytotoxic effect of Schiff base copper complexes, cell viability study was done with the MTT reduction assay. Hep-2 cells were seeded in a 96-well plate at the density of 5x10³ cells/well. The cells were allowed to attach and were grown in 96-well plate for 24 h, in 200 μ L of Eagle's minimal essential medium (EMEM) with 10% Fetal Bovine Serum (FBS) [38]. After that the media was removed and replaced with suspension of various concentrations of Schiff base copper complexes (10 to 100 mg/mL) (minimum 4 wells seeded with each concentration), the cells were incubated for 48 h. After the addition of (MTT), 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (10 mL, 5 mg/mL), the cells were incubated at 37° C for another 4 h. The medium was then removed, and 200 μ L of DMSO was added to each well. Optical density of the formazan product was read at 620 nm using multi well spectrophotometer. The results were given as mean of four independent experiments. OD value was subjected to sort out percentage of viability by using the following formula

$$\text{Percentage of cell viability} = \frac{\text{OD value of the treated sample}}{\text{OD value of the control sample}} \times 100$$

Molecular Docking Study

Human-DNA-Topo-I complex (PDBID: 1SC7)] and Dengue NS3 protease-helicase bi-functional enzyme (PDB ID: 2VBC) were obtained from Protein Data Bank. Metal complexes were converted into PDB format using Mercury software [39]. The 'receptor' (DNA/Dengue) and 'ligand' (metal complexes) files were docked using Auto Dock Tools. The heteroatoms including water molecules and excess ligand were removed and also polar hydrogen atoms and Kollman charges were added to the receptor molecule to investigate the binding nature. In the binding was enclosed in a box which having number of grid points in (x) x (y) x (z) directions of 40 x 40 x 40 and a grid spacing of 0.35 Å. Docking studies were conducted using Auto Dock Tools (ADT) version 1.5.4 and finalized by Auto Dock vina programme. The docked structures were exposed using Discovery studio software.

3. RESULTS & DISCUSSION

¹H-NMR Spectral Studies and Electron Ionization Mass Spectral analysis

The ¹H NMR spectra of the ligand L is provided in the Fig. 1. The appearance of signal at 8.477 ppm is attributed to azomethine group [40] and the signal for amido proton appeared at 8.513 ppm [41]. The atomic protons of the ligand appeared in the range 7.2 – 7.9 ppm, which confirms the formation of the ligand. The EI mass spectra of ligand L is given in Fig. 2 which has the molecular ion M⁺ peak at 492.9514 corresponding to the molecular weight of the ligand L. The peaks at 474.9596, 457.0669, 373.0051, 353.0565, 248.9471, 141.9279, 114.0282, 103.04 and 76.1198 corresponds to various fragments of ligand L.





Fig. 1: ¹H NMR spectrum of L



→ 2E_g and $^2B_{1g} \rightarrow ^2B_{2g}$ transitions. This further stands as the evidence for the square planar geometry of d^9 Cu(II) system [51].

EPR spectral analysis

The EPR spectra of copper complex provide information of importance in studying the metal ion environment. The copper complex $Cu(L)_2$ exhibits an isotropic signal, without any hyperfine splitting, with $g_{iso} = 2.104$ as shown in the Fig. 5. The g value obtained in the present study when compared to the g value of a free electron, 2.0023, indicate an increase of the covalent nature of the bonding between the metal ion and the ligand molecule [52]. Isotropic lines usually results due to occupancy of the unpaired electron in a degenerate orbital in square planar geometry.




Fig. 5: EPR Spectrum of $Cu(L)_2$

Cytotoxicity screening

There are several methods for assessing the carcinogenic or mutagenic properties of the given chemical structure. The method followed here is a

bacteria strain based assay, which includes *E. coli* AB1157 which is a wild-type strain proficient in DNA damage repair. The bacterial strain is incubated with the compounds of interest for the analysis of any associated lethal effects. On incubation with the compounds, any free radical generation by the compounds lead to the lethality of the cells [53]. This cytotoxic potentiality of the compound will be displayed in terms of zone of inhibition. The cytotoxic screening analysis of the complexes showed that copper complex alone exhibit excellent cytotoxicity screening effects at MIC value of 0.25 mg.

Table 1. Zone of inhibition (mm) for Cytotoxicity Screening analysis

Compounds	0.125 mg	0.25 mg	0.5 mg	1.0 mg	2.0 mg	MIC (mg)
$Co(L)_2$	0	0	3	7	10	0.5
$Ni(L)_2$	0	0	0	0	5	2.0
$Cu(L)_2$	0	2	4	5	10	0.25
Stannous chloride	0	0	3	8	15	0.5

DNA binding studies using electronic spectra

The interactions of metal complexes with DNA are of interest in order for the development of chemotherapeutic agents. Electronic absorption spectroscopy is one of the most useful techniques for DNA binding studies of metal complexes. The interactions of copper complex $Cu(L)_2$ with CT-DNA were investigated by UV-Vis absorption titrations and is represented in Fig. 6.




Fig. 6: Cytotoxicity Screening analysis of complexes

Upon addition of increasing amount of CT-DNA from 0 – 200 μL , a significant “hyperchromic” effect of the intra ligand bands at 257.8 – 300 nm was accompanied by a red shift of 2 – 7 nm, indicative of the breakage of the DNA helix [54,55]. There is no appreciable change in the charge transfer band. As the concentration of the DNA was increased, the absorption bands of the copper complex initially showed hyperchromism, but on further increasing concentration, hyperchromism with blue shift is obtained (fig. 7).




Fig. 7: DNA Binding interaction of $\text{Cu}(\text{L})_2$

The invitro cytotoxicity of the copper complex $\text{Cu}(\text{L})_2$ was evaluated against Larynx Cancer Cells Hep_2 and given in fig. 8. The IC_{50} (the concentrations that inhibited in 50% the cellular proliferation) of the studied complex is $40\mu\text{g}$.

Molecular docking with human DNA topoisomerase I

Human DNA topoisomerase I and II were the selective targeting area for synthesizing the anticancer drug [56]. The molecular docking of Cobalt(II),

Nickel(II) and Copper(II) complexes were performed to determine the value of binding affinity and the selected binding residue which along with the sterically suitable conformations. The low value of the binding energy shows the more effective binding affinity between the ‘receptor’ and the ‘ligand’ molecules. The various conformations of docked molecular complexes were analyzed in terms of binding energy, hydrogen bonding and hydrophobic interaction between receptors and the acceptor. More negative value of the relative binding values suggests that the interaction between the DNA and ligand is so strong, due to the extended aromatic ring. Phenyl ring has higher free binding energy which gives a better binding affinity value comparing with a compound containing bipyridyl ring. From these work mononuclear complexes gives better results towards the HDNA. The binding energy values of the $\text{Co}(\text{II})$, $\text{Ni}(\text{II})$ and $\text{Cu}(\text{II})$ complexes respectively are -11.6, -6.9 and -10.4 kcal mol^{-1} towards human DNA topoisomerase I. This shows that Cobalt(II) and Copper(II) molecule easily binds with the DNA helix and the Nickel(II) complex preferred to bind with the outermost protein’s amino acid residue. The docked images were shown in Fig. 9.

NS3 protease-helicase (dengue virus) is a very important target area which should be docked. Cobalt(II), Nickel(II) and Copper(II) complexes exhibit very low binding energy value, it means that complexes having very high binding affinity towards the NS3 protease-helicase. The distance between the selected receptor to targeted molecule is also low. The binding energy values of the $\text{Co}(\text{II})$, $\text{Ni}(\text{II})$ and $\text{Cu}(\text{II})$ complexes are -12.7, -12.9 and -12.6 kcal mol^{-1} respectively. The binding interactions were shown in fig.10. From the theoretical point of view these complexes are consider to be a good anti-dengue drug.




Fig. 8: Anti-Tumor activity of $\text{Cu}(\text{L})_2$

[illegible]

17

4. CONCLUSION

A new Schiff base ligand L which contains several phenyl rings and its cobalt(II), nickel(II) and copper(II) complexes were synthesized. The formation of the ligand and its metal complexes were confirmed by various spectroscopic techniques. The biological applications of the synthesized complexes were carried out by Cytotoxic screening analysis, DNA binding Ability by using electronic spectra and Anti-Tumor activity by MTT assay. The results obtained showed that among the synthesized complexes copper complex has potential biological activity. The docking studies were carried out using synthesized metal complexes with human DNA topoisomerase I (PDB: 1SC7) and Dengue NS3 protease-helicase bi-functional enzyme (PDB ID: 2VBC) using Auto Dock vina and Discovery studio software. The binding energy values of the Co(II), Ni(II) and Cu(II) complexes showed that these types of compounds can be act as an potential Anti-Dengue and Anticancer agent

ACKNOWLEDGEMENTS

The authors are thankful to Indian Institute of Technology Chennai for providing their help in taking EI-mass and EPR spectra and also Dr. Gnanavel for his support to do this work.

REFERENCES

- Abbaspour, A., Esmailbeig, A. R., Jarrahpour, A. A., Khajeh, B., Kia, R. Talanta, 58, 397–403(2002).
doi:10.1016/S0039-9140(02)00290-4.
- Abdulghani, A. J. and Mohuee, S. K., ‘Synthesis and Characterization of New Schiff Bases Derived from Ceftriaxone Sodium with α H- Indole-2, 3-Dione (Isatin) and α -Acetyl Indoline-2, 3-Dione (N-Acetylisatin) and their Platinum(IV) complexes’, Journal of Chemical Biological and physical Sciences, 6(2), 579-595(2016).
- Alagesan, M., N. S. P. Bhuvanesh, N. Dharmaraj, An investigation on new ruthenium (II) hydrazone complexes as anticancer agents and their interaction with biomolecules, Dalton Trans., 43, 6087-6099(2014).
doi:10.1039/C3DT51949J
- Aoyama, Y., Konoike, T., Kanda, A., Naya, N. and Nakajima, M., Total synthesis of human chymase inhibitor methylinderone and structure–activity relationships of its derivatives, Bioorg. Med. Chem. Lett., 11(13), 1695–1697(2001).
doi: 10.1016/S0960-894X(01)00265-7
- Aoyama, Y., Uenaka, M., Kii, M., Tanaka, M., Konoike, T., Hayasaki - Kajiwar Y., Naya N. and Nakajima, M., Design, synthesis and pharmacological evaluation of 3-benzylazetidine-2-one-based human chymase inhibitors, Bioorg. Med. Chem., 9(11), 3065–3075(2001).
doi:10.1016/S0968-0896(01)00209-7.
- Aoyama Y., Uenaka M., Konoike T., Hayasaki-Kajiwar Y., Naya N., Nakajima M., Inhibition of serine proteases: activity of 1, 3-Diazetidine-2, 4-diones, Bioorg. Med. Chem. Lett., 11(13), 1691–1694(2001).
doi:10.1016/S0960-894X(01)00264-5.
- Aoyama, Y., Uenaka, M., Konoike, T., Iso, Y., Nishitani, Y., Kanda, A., Naya, N. and Nakajima M., Synthesis and structure-activity relationships of a new class of 1-oxacephem-based human chymase inhibitors, Bioorg. Med. Chem. Lett., 10(21), 2397–2401(2000).
doi:10.1016/S0960-894X(00)00488-1.
- Asadi, Mozaffar, Elham Safaei, Bijan Ranjbar & Leila Hasani, Thermodynamic and spectroscopic study on the binding of cationic Zn(II) and Co(II) tetrapyrrolineporphyrins to calf thymus DNA: the role of the central metal in binding parameters, New Journal of Chemistry, 28(10), 1227-1234(2004).
doi.org/10.1039/B404068F
- Bas, D. C., Rogers, D. M. and Jensen, J. H., Very fast prediction and rationalization of pKa values for protein-ligand complexes, Proteins Struct. Funct. Genet., 73(3), 765–783(2008).
doi: 10.1002/prot.22102.
- Bellamy, L. J., The Infrared Spectra of Complex Molecules, Chapman and Hall, London(1978).
- Chen L. S. and Sue C. Cummings, Synthesis and characterization of cobalt (II) and some nickel (II) complexes with N, N'-ethylenebis (pX-benzoylacetone iminato) and N, N'-ethylenebis (pX-benzoylmonothioacetone iminato) ligands, J. Inorg. Chem., 17(9), 2358-2361(1978).
doi.org/10.1021/ic50187a005
- Del Paggio, Alan A & David R. McMillin 1983, ‘Substituent effects and the photoluminescence of Cu (PPh₃)₂ (NN)⁺ systems’, Journal of Inorganic Chemistry, vol. 22, no. 4, pp. 691-692.
- Geary, W. J., Coord. Chem. Rev., 7 (1971)
- Greco M.N., Hawkins M.J., Powell E.T., Almond H.R., Garavilla L., Hall J., Minor L.K., Wang Y., Corcoran T.W., Cera E.D., Cantwell A.M., Savvides S.N., Damiano B.P., Maryanoff B.E. (2007) Discovery of potent, selective, orally active, nonpeptide inhibitors of human mast cell chymase. J Med Chem;50:1727–1730.
- Greenwood, J. R., D. Calkins, A.P. Sullivan, J.C. Shelley, Towards the comprehensive, rapid, and accurate prediction of the favorable tautomeric states of drug-like molecules in aqueous

- solution, J. Comput. Aided Mol. Des.24 (2010) 591–604
- Greenwood, Jeremy R, David Calkins, Arron P. Sullivan & John C. Shelley 2010, 'Towards the comprehensive, rapid and accurate prediction of the favourable tautomeric states of drug-like molecules in aqueous solution', Journal of Computer-aided Molecular Design, vol. 24, no. 6-7, pp. 591-604.
- Hayashi Y., Iijima K., Katada J., Kiso Y. (2000) Structure-activity relationship studies of chloromethyl ketone derivatives for selective human chymase inhibitors. Bioorg Med ChemLett;10:199- 201.
- Herebian, Diran & William S. Sheldrick 2002, 'Synthesis and DNA binding properties of bioorganometallic (η 5-pentamethylcyclopentadienyl) iridium (III) complexes of the type [η 5-C 5 Me 5) Ir (Aa)(dppz)] n+ (dppz=dipyrido [3, 2-a' 2', 3'-c] phenazine, n=1-3), with S-coordinated amino acids (Aa) or peptides', Journal of the Chemical Society, Dalton Transactions, vol. 6, pp. 966-974.
- Iijima K., Katada J., Hayashi Y. (1999) Symmetrical anhydridetype serine protease inhibitors:structure-activity relationship studies of human chymase inhibitors. Bioorg Med ChemLett;9:413–418.
- Iijima K., Katada J., Yasuda E., Uno I., Hayashi Y. (1999) N-[2,2-Dimethyl-3-(N-(4-Cyanobenzoyl)Amino)Nonanoyl]-L-Phenylalanine ethyl ester as a stable ester-type inhibitor of chymotrypsin-like serine proteases: structural requirements for potent inhibition of a-chymotrypsin. J Med Chem;42:312–323.
- Inas Ahmed Mohammed Saif 2017,, 'A mini Review of Complexes of Schiff Base and their Biological Activities', Journal of Medicinal Chemistry and Drug Discovery, vol. 2, no. 3, pp. 93-107.
- Irwin, J. J., B.K. Shoichet, ZINC—a free database of commercially availablecompounds for virtual screening, J. Chem. Inf. Model. 45 (2005) 177–182
- Jamieson, E. R., S. J. Lippard, Chem. Rev. 1999, 99, 2467.
- Kemp & William 1991, 'Energy and the Electromagnetic Spectrum' in Organic Spectroscopy, pp. 1-17, Palgrave, London.
- Khalil, S. M. E., H. S. Seleem, B. A. El-Shetary, M. Shebl M., J. Coord. Chem., 55(8) (2002) p. 883. <http://dx.doi.org/10.1080/0095897022000002213>.
- Krishnaraj C, Muthukumaran P, Ramachandran R, Balakumaran MD & Kalaichelvan PT 2014, 'Acalypha indica Linn: biogenic synthesis of silver and gold nanoparticles and their cytotoxic effects against MDA-MB-231, human breast cancer cells', Journal of Biotechnology Reports, vol. 4, pp. 42-49.
- Kriza, A., A.Reiss, S.Florea, T. Caproiu, J.Indian chem. Soc. 77 (2000) p. 207-208.
- Lemos PRG, On Terra Junior, Anantea ML, Nunes AS & Soares JRO 2011, 'Evaluation of the effects of an aqueous extract of Schinus terebinthifolius involving wild bacterial culture of Escherichia coli', Asian Journal of Pharmaceutical and Health, vol. 1, pp. 86-88.
- Leung, M. H. M., T. Harada, T. W. Kee, Curr. Pharm. Des. 2013, 19, 2070
- Li, H., A.D. Robertson, J.H. Jensen, Very fast empirical prediction andrationalization of protein pKavalues, Proteins Struct. Funct. Genet. 61 (2005)704–721
- Liu, H. K., P. J. Sadler, Acc. Chem. Res. 2011, 44, 349
- Madhavi Sastry, G., M. Adzhigirey, T. Day, R. Annabhimoju, W. Sherman,Protein and ligand preparation: parameters, protocols, and influence onvirtual screening enrichments, J. Comput. Aided Mol. Des. 27 (2013) 221–234
- Mjos, K. D., C. Orvig, Chem. Rev. 2014, 114, 4540
- Murphy, F. A., B.W.J. Mahy, M.H.V. van Regenmortel, Epidemiology of humanand animal viral diseases, Encycl. Virol. (2008) 140–148.
- Natarajan C, Tharmaraj P & Murugesan R 1992, 'In situ synthesis and spectroscopic studies of copper (II) and nickel (II) complexes of 1-hydroxy-2-naphthylstyrylketoneimines', Journal of Coordination Chemistry, vol. 26, no. 3, pp. 205-213.
- Niwata S., Fukami H., Sumida M., Ito A., Kakutani S., Saitoh M., Suzuki K. et al. (1997) Substituted 3-(Phenylsulfonyl)-1- phenylimidazolidine-2,4-dione derivatives as novel nonpeptide inhibitors of human heart chymase. J Med Chem;40:2156–2163.
- Noble, C. G., C.C. Seh, A.T. Chao, P.Y. Shi, Ligand-bound structures of the denguevirus protease reveal the active conformation, J. Virol. 86 (2012) 438–446
- Ortiz, Bertha & Su-Moon Park 2000, 'Electrochemical and spectro electrochemical studies of cobalt salen and salophen as oxygen reduction catalysts', Bulletin-Korean Chemical Society, vol. 21, no. 4, pp. 405-411.
- Patel KV & Bhattacharya PK 1982, 'Studies in Binuclear Copper (II) Complexes of Tetradentate Schiff-Bases Involving Greater π -Delocalization', Indian Journal of Chemistry Section A-Inorganic Bio-Inorganic Physical Theoretical & Analytical Chemistry, vol. 21, no. 12, pp. 1110-1112.
- Pommier Y *Nat. Rev. Cancer.*, **2006**, 6(10), 789-802.
- Raman N & Sobha S 2012, 'Exploring the DNA binding mode of transition metal based biologically active compounds', Spectro Chimica Acta Part A: Molecular and Biomolecular Spectroscopy, vol. 85, no. 1, pp. 223-234.
- Raman N, Dhavethu Raja J & Sakthivel A 2007, 'Synthesis, spectral characterization of Schiff base transition metal complexes: DNA cleavage and

- antimicrobial activity studies', *Journal of Chemical Sciences*, vol. 119, no. 4, pp. 303-310.
- Ramesh R & Maheswaran S 2003, 'Synthesis, spectra, dioxygen affinity and antifungal activity of Ru (III) Schiff base complexes', *Journal of Inorganic Biochemistry*, vol. 96, no. 4, pp. 457-462.
- Schiff HS 1864, 'Sur quelques derives pheniques des aldehydes', *Annali Di Chimica*, vol. 131. pp. 118.
- Schiff, H. *Ann. Chem.* 1869, 150, p. 193.
- Schiff, H. *Ann. Chem.* 1869, 151, p. 186.
- Schiff, H. *Ann. Chem. Suppl.* 1864, 3, p. 343.
- Seleem, H. S., *Ann. Chim.*, 93 (2003) p. 305.
- Shakir, Mohammed, Omar SM Nasman, Ajax K. Mohamed & Saji P. Varkey 1996, 'Transition metal complexes of 13-14-membered tetraazamacrocycles: Synthesis and Characterization', *Polyhedron*, vol. 15, no. 8, pp. 1283-1287.
- Shao, J., Z. Y. Ma, A. Li, Y. H. Liu, C. Z. Xie, Z. Y. Qiang, J. Y. Xu, *J. Inorg. Biochem.* 2014, 136, 13.
- Sigman, D. S., A. Mazumder, D. M. Perrin, *Chem. Rev.* 1993, 93, 229.
- Singh DP, Vandna Malik, Krishan Kumar, Chetan Sharma & Aneja KR 2010, 'Macrocyclic metal complexes derived from 2, 6-diaminopyridine and isatin with their antibacterial and spectroscopic studies', *Spectro Chimica Acta Part A: Molecular and Biomolecular Spectroscopy*, vol. 76, no. 1, pp. 45-49.
- Tadele KT 2017, 'Antioxidant Activity of Schiff Bases and Their Metal Complexes: A Recent Review', *Journal of Pharmaceutical and Medicinal Research*, vol. 3, no. 1, pp. 73-77.
- Thomas, Marykutty, Muraleedharan Nair MK & Radhakrishnan PK 1995, 'Rare earth iodide complexes of 4-(2', 4'-dihydroxyphenylazo) antipyrine', *Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry*, vol. 25, no. 3, pp. 471-479.
- Vogel, A., *Text Book of Quantitative Inorganic Analysis* third Edition, ELBS London 1969.
- Wichapong, K., A. Nueangaudom, S. Pianwanit, W. Sippl, S. Kokpol, Identification of potential hit compounds for Dengue virus NS2B/NS3protease inhibitors by combining virtual screening and binding free energy calculations, *Trop. Biomed.* 30 (2013) 388–408.